

On the Ricci curvature of normal metrics on Biquotients

Lorenz J. Schwachhöfer*

February 21, 2007

Abstract

We show that any normal metric on a closed biquotient with finite fundamental group has positive Ricci curvature.

1 Introduction

One of the classical problems in differential geometry is the investigation of closed manifolds which admit Riemannian metrics with given lower bounds for the sectional or the Ricci curvature. As a sample question, one may ask for manifolds with a metric of both nonnegative sectional and positive Ricci curvature. Note that such a manifold has necessarily a finite fundamental group by the Bonnet-Myers theorem.

By a recent result of Böhm and Wilking ([BW]), a closed manifold with finite fundamental group which carries a metric of nonnegative sectional curvature also admits metrics of positive Ricci curvature. These metrics are obtained by a deformation of the original metric using the Ricci flow, so we may assume in addition that these metrics have *almost* nonnegative sectional curvature.

In general, however, manifolds which have nonnegative sectional and positive Ricci curvature simultaneously seem to be hard to come by. One class of such manifolds are compact homogeneous spaces G/H with a *normal* metric, i.e., a metric for which the canonical submersion $G \rightarrow G/H$ is Riemannian where G is equipped with a biinvariant metric. Such a metric always has nonnegative sectional curvature by O'Neill's formula, and it is not hard to see that it has positive Ricci curvature as long as G/H has finite fundamental group ([N]).

Another class of manifolds of nonnegative sectional curvature are closed biquotients $G//H$ for $H \subset G \times G$ with a *normal* metric, by which we mean - in analogy to the homogeneous case - a metric for which the canonical submersion $G \rightarrow G//H$ is Riemannian, G being equipped with a biinvariant metric. Normal metrics are in some sense the most canonical metrics on closed biquotients, so the question under which circumstances such a metric has positive Ricci curvature is very natural.

Surprisingly, not much was known about this question up to now. In [F], this positivity was shown under very restrictive circumstances; in [ST], we showed that a normal metric on a biquotient with finite fundamental group has positive Ricci curvature on a dense open subset. The aim of the present article is to resolve this question completely.

*Research supported by the Schwerpunktprogramm Differentialgeometrie of the Deutsche Forschungsgesellschaft

Main Theorem *Let $M := G//H$ be a biquotient of a compact Lie group G , equipped with a normal metric. If M has finite fundamental group then M has positive Ricci curvature.*

Thus, any closed biquotient with finite fundamental group carries a metric of both nonnegative sectional and positive Ricci curvature. In general, we show that a closed biquotient with a normal metric is finitely isometrically covered by $M' \times T^k$, where M' is a biquotient with a normal metric of positive Ricci curvature and T^k is a flat torus.

It is my pleasure to thank the Max-Planck-Institute for Mathematics in the Sciences in Leipzig and the Université Libre de Bruxelles for their hospitality during the preparation of parts of this work.

2 Normal metrics on Biquotients

Let G be a compact connected Lie group with Lie algebra $\mathfrak{g}$, and let $H \subset G \times G$ be a closed subgroup for which the action of H on G given by

$$(h_1, h_2) \cdot g := h_1 g h_2^{-1}$$

is free and hence the quotient $G//H$ is a manifold, called the *biquotient of G by H* .

Let Q be an Ad_G -invariant inner product on $\mathfrak{g}$ which induces a biinvariant Riemannian metric of nonnegative sectional curvature on G . Since the action of H on G preserves Q , there is a unique Riemannian metric g_Q on $G//H$ for which the projection

$$\pi : G \longrightarrow G//H$$

becomes a Riemannian submersion. Evidently, g_Q has nonnegative sectional curvature by O'Neill's formula.

Definition 2.1 *A metric g_Q on $G//H$ induced by the Ad_G -invariant inner product Q on $\mathfrak{g}$ as above is called normal.*

If $H = L \times K \subset G \times G$ then H acts freely on G iff $Ad_g L \cap K = \{e\}$ for all $g \in G$. In this case, we denote the biquotient $G//H$ also by $L \backslash G / K$. In fact, *any* biquotient can be considered to be of this form with G/K being a *symmetric space*. Namely, there is a canonical diffeomorphism

$$G//H \longleftrightarrow H \backslash (G \times G) / \Delta G. \quad (1)$$

If we consider the normal metric $g_{Q \oplus Q}$ on $H \backslash (G \times G) / \Delta G$, then one sees easily that the diffeomorphism in (1) becomes an isometry. Therefore, since $(G \times G) / \Delta G$ is a symmetric space, we shall from now on always consider biquotients of the form

$$M = H \backslash G / K, \quad \text{where } G/K \text{ is a symmetric space,} \quad (2)$$

and we have the corresponding involution τ , given by

$$\tau : \mathfrak{g} \rightarrow \mathfrak{g}, \quad \tau|_{\mathfrak{k}} = Id_{\mathfrak{k}}, \quad \tau|_{\mathfrak{k}^\perp} = -Id_{\mathfrak{k}^\perp}. \quad (3)$$

For $g \in G$, we define the Q -orthogonal decomposition

$$\mathfrak{g} = \mathcal{V}_g \oplus \mathcal{H}_g, \quad \text{where } \mathcal{V}_g := Ad_{g^{-1}}(\mathfrak{h}) \oplus \mathfrak{k}, \quad (4)$$

with the Lie algebras $\mathfrak{h}$ and $\mathfrak{k}$ of $H \subset G$ and $K \subset G$, respectively. Note that $dL_g(\mathcal{V}_g) = \ker(d\pi_g)$, so that $T_g G = dL_g(\mathcal{V}_g) \oplus dL_g(\mathcal{H}_g)$ is the canonical decomposition of TG into the vertical and horizontal subspace w.r.t. the Riemannian submersion $\pi : G \rightarrow G//H$. In particular, the dimensions of $\mathcal{V}_g$ and $\mathcal{H}_g$ are independent of g .

3 Biquotients with a flat point

As a first step towards our main result, we investigate biquotients with flat points, i.e., points where the curvature vanishes entirely. Since $\pi : G \rightarrow G//H$ is a Riemannian submersion, hence $Sec_{\pi(g)}^{G//H}(\underline{v} \wedge \underline{w}) \geq Sec_g^G(v \wedge w) = \frac{1}{4} \|[v, w]\|^2$, where $v, w \in \mathcal{H}_g$ are horizontal lifts of $\underline{v}$ and $\underline{w}$, respectively, it follows that $\pi(g) \in G//H$ is a flat point only if $[\mathcal{H}_g, \mathcal{H}_g] = 0$. In this case, we can prove the following

Theorem 3.1 *Let $G//H$ be a biquotient with a normal metric. If $G//H$ has a flat point, i.e., if there is a $g \in G$ with $[\mathcal{H}_g, \mathcal{H}_g] = 0$, then $\mathcal{H}_g \subset \mathfrak{z}(\mathfrak{g})$. In this case, $G//H$ is finitely covered by a flat torus.*

Proof. We assume w.l.o.g. that $g = e$. After passing to a finite cover, we further assume that $G = G_s \times Z$, where G_s is semi-simple and simply connected and $Z = Z(G)$ is the identity component of the center of G . Moreover, we may assume that $H \subset G \times G$ is connected. We have an orthogonal splitting of the Lie algebra $\mathfrak{g} = \mathfrak{g}_s \oplus \mathfrak{z}(\mathfrak{g})$, and we let $T \subset \overline{\exp \mathcal{H}_e} \subset G$ be a torus with Lie algebra $\mathfrak{t}$ such that $\mathfrak{g} = \mathcal{V}_e \oplus \mathfrak{t}$ as a vector space. In particular, $\dim T = \dim \mathcal{H}_e$. (We use T since the abelian subgroup $\exp \mathcal{H}_e \subset G$ may be not closed). Now consider the map $p : T \rightarrow G//H$ which makes the following diagram commute.

$$\begin{array}{ccc} & & G \\ & \nearrow & \downarrow \pi \\ T & & G//H \\ & \searrow p & \end{array} \quad (5)$$

Let $g \in T$. Since $T \subset \overline{\exp(\mathcal{H}_e)}$ and the latter is abelian, we have

$$Q(Ad_{g^{-1}}(\mathfrak{h}), \mathcal{H}_e) = Q(\mathfrak{h}, Ad_g(\mathcal{H}_e)) = Q(\mathfrak{h}, \mathcal{H}_e) = 0, \quad \text{and} \quad Q(\mathfrak{k}, \mathcal{H}_e) = 0.$$

Thus, $Q(\mathcal{V}_g, \mathcal{H}_e) = 0$, which implies that $\mathcal{V}_g = \mathcal{V}_e$ for all $g \in T$. In particular, $\mathfrak{t}$ is transversal to $\mathcal{V}_g$ for all $g \in T$. As $\mathcal{V}_g$ is by definition the kernel of the differential $d\pi_g : T_g G \rightarrow T_{p(g)} G//H$ for $g \in T$, it follows that $dp_g : dL_g(\mathfrak{t}) \hookrightarrow T_g G \rightarrow d_{p(g)} G//H$ is an isomorphism, hence $p : T \rightarrow G//H$ is a finite covering map, hence the induced map $p_* : \pi_1(T) \rightarrow \pi_1(G//H)$ is injective with index $[\pi_1(G//H) : p_*(\pi_1(T))] < \infty$.

Since $G = G_s \times Z$ with G_s simply connected, it follows that the projections $pr_Z : G \rightarrow Z$ induces an isomorphism of fundamental groups, hence from the homotopy exact sequence

$$\begin{array}{ccccc} & \pi_1(T) & & & \\ & \downarrow (pr_Z)_* & \searrow p_* & & \\ \pi_1(H) & \xrightarrow{(pr_Z)_*} & \pi_1(Z) & \xrightarrow{\pi_*} & \pi_1(G//H) \longrightarrow 0 \end{array}$$

and from $[\pi_1(G//H) : p_*(\pi_1(T))] < \infty$ we conclude that $\dim pr_{\mathfrak{z}(\mathfrak{g})}(\mathfrak{t}) = \dim \mathfrak{t}$ and that $\mathfrak{z}(\mathfrak{g}) = pr_{\mathfrak{z}(\mathfrak{g})}(\mathcal{V}_e) \oplus pr_{\mathfrak{z}(\mathfrak{g})}(\mathfrak{t})$. Then, for dimensional reasons, it follows that $\mathcal{V}_e = pr_{\mathfrak{z}(\mathfrak{g})}(\mathcal{V}_e) \oplus \mathfrak{g}_s$ and therefore, $\mathcal{H}_e \subset \mathfrak{g}_s^\perp = \mathfrak{z}(\mathfrak{g})$. $\blacksquare$

4 Biquotients with Ricci-flat directions

Let us suppose that at a point $p \in G//H$ the Ricci curvature is not positive, i.e., there are tangent vectors $0 \neq \underline{v} \in T_p G//H$ with $Sec_{\pi(g)}^{G//H}(\underline{v} \wedge \underline{w}) = 0$ for all $\underline{w} \in T_p G//H$. W.l.o.g. we assume that $p = \pi(e)$. As in the preceding section, O'Neill's formula then implies that $[v, \mathcal{H}_e] = 0$ where $v \in \mathcal{H}_e$ is the horizontal lift of $\underline{v}$. To simplify our notation we shall from now on omit the subscript and thus let $\mathcal{V} := \mathcal{V}_e = \mathfrak{h} \oplus \mathfrak{k}$ and $\mathcal{H} := \mathcal{H}_e = \mathcal{V}^\perp$. Thus, we define the space of Ricci flat directions at $\pi(e)$

$$\mathcal{F} := \{v \in \mathcal{H} \mid [v, \mathcal{H}] = 0\} \subset \mathcal{H} \quad (6)$$

and assume that $\mathcal{F} \neq 0$.

Proposition 4.1 *Let $M := H \backslash G / K$ be a normal biquotient as in (2), and suppose that $\mathcal{F} \subset \mathfrak{z}(\mathfrak{g})$. Then M is finitely isometrically covered by $M' \times T^k$, where M' is a biquotient with finite fundamental group and a normal metric, and T^k is a flat torus of dimension $k = \dim \mathcal{F}$.*

Proof. After replacing G by a finite cover if necessary, we may assume that $G = G_s \times Z$ where G_s is semisimple and Z is the identity component of the center with Lie algebra $\mathfrak{z}(\mathfrak{g})$. Moreover, after replacing M by a finite isometric cover, we may assume that $H, K \subset G$ are connected. Let $pr_Z : G = G_s \times Z \rightarrow Z$ be the canonical projection.

Note that the map $j : H \times K \rightarrow Z$, $(h, k) \mapsto pr_Z(h) \cdot pr_Z(k)$ is a homomorphism, hence its image $j(H \times K) \subset Z$ is a torus. Also, by hypothesis, it follows that $\mathcal{F} = \mathfrak{z}(\mathfrak{g}) \cap \mathcal{H}$, so that $\mathfrak{z}(\mathfrak{g}) = pr_{\mathfrak{z}(\mathfrak{g})}(\mathfrak{h} \oplus \mathfrak{k}) \oplus \mathcal{F}$.

Let $G' := G_s \times j(H \times K)$. Then evidently, $G' \triangleleft G$, and $G \cong G' \times T^k$, where $T^k \subset Z$ is a torus with Lie algebra $\mathfrak{t} \subset \mathfrak{z}(\mathfrak{g})$ of dimension $k = \dim(\mathcal{F})$.

Choose a linear map $p : \mathfrak{g} \rightarrow \mathfrak{g}$ such that $p|_{\mathfrak{g}'} = Id_{\mathfrak{g}'}$ and $p(\mathfrak{t}) = \mathcal{F}$, and define the inner product $\tilde{Q} := p^*(Q)$. Since $\tilde{Q}|_{\mathfrak{g}'} = Q|_{\mathfrak{g}'}$ and $\tilde{Q}(\mathfrak{g}_s, \mathfrak{z}(\mathfrak{g})) = 0$, it follows that $\tilde{Q}$ is biinvariant as well. Note that $H \times K \subset G'$, hence we have a canonical isometry

$$(H \backslash G / K, g_Q) \longleftrightarrow ((H \backslash G' / K) \times T^k, g_{\tilde{Q}}),$$

where the latter is a product metric, and the metric on the torus factor is flat. Also, by construction, $M' := ((H \backslash G' / K), g_{\tilde{Q}}) = ((H \backslash G' / K), g_Q)$ is a biquotient with a normal metric which has positive Ricci curvature at $\pi(e)$. Therefore, by the deformation result of Ehrlich [E], M' admits a metric of positive Ricci curvature and hence has finite fundamental group. $\blacksquare$

Definition 4.2 *Let $M := H \backslash G / K$ be a normal biquotient as in (2). We say that M has non-central flats if $\mathcal{F} \not\subset \mathfrak{z}(\mathfrak{g})$. If M has non-central flats, then we call M minimal if for any normal biquotient $M' = H' \backslash G' / K'$ with non-central flats with $\dim M' \leq \dim M$ and $\dim G' \leq \dim G$ we have $\dim M' = \dim M$ and $\dim G' = \dim G$.*

Thus, our main theorem will follow if we can show that there do not exist normal biquotients with non-central flats which shall be our goal for the remainder of this paper and will be achieved in Theorem 4.14. Indeed, once this is shown it will follow from Proposition 4.1 that a biquotient with a normal metric with Ricci flat directions has infinite fundamental group.

Proposition 4.3 *Suppose that $M = H \backslash G / K$ is a minimal biquotient with non-central flats. Then G / K is a compact irreducible symmetric space.*

Proof. Suppose that $\mathfrak{g} = \mathfrak{g}_1 \oplus \mathfrak{g}_2$ is a τ -invariant decomposition where $\tau : \mathfrak{g} \rightarrow \mathfrak{g}$ denotes the symmetric involution from (3). After passing to a covering if necessary, we may assume w.l.o.g. that $G / K = G_1 / K_1 \times G_2 / K_2$ where $G = G_1 \times G_2$ and $K_i := K \cap G_i$. Then $\mathfrak{k} = (\mathfrak{k} \cap \mathfrak{g}_1) \oplus (\mathfrak{k} \cap \mathfrak{g}_2) =: \mathfrak{k}_1 \oplus \mathfrak{k}_2$. Let $pr_i : \mathfrak{g} \rightarrow \mathfrak{g}_i$ be the canonical projection. Let $\mathcal{H}_i := pr_i(\mathcal{H})$ and $\mathcal{F}_i := pr_i(\mathcal{F}) \subset \mathcal{H}_i \subset \mathfrak{k}_i^\perp$. If $\mathcal{F}_i \subset \mathfrak{z}(\mathfrak{g}_i)$ for $i = 1, 2$ then we would have $\mathcal{F} \subset \mathfrak{z}(\mathfrak{g})$ which we assumed *not* to be the case, thus we may assume w.l.o.g. that $\mathcal{F}_1 \not\subset \mathfrak{z}(\mathfrak{g}_1)$.

Now consider the biquotient $M' := (H \cap G_1) \backslash G_1 / K_1$. Its horizontal space is $\mathfrak{k}_1^\perp \cap (\mathfrak{h} \cap \mathfrak{g}_1)^\perp = \mathcal{H}_1$, hence $\mathcal{F}_1 \subset \mathcal{H}_1$ is the space of flat directions of M' . Since $\mathcal{F}_1 \not\subset \mathfrak{z}(\mathfrak{g}_1)$, it follows that M' has non-central flats, and $\dim M' = \dim \mathcal{H}_1 \leq \dim \mathcal{H} = \dim M$. But $\dim G_1 \leq \dim G$, hence the minimality of M implies that $\dim G_1 = \dim G$, i.e., $G_2 = 1$. $\blacksquare$

The following simple facts will be useful later on.

Lemma 4.4 *Let $\mathfrak{g}$ be a Lie algebra with a biinvariant inner product Q , and let $L \subset \mathfrak{g}$ be a linear subspace. Let $L_1 := [L, \mathfrak{g}]$. Then the ideal generated by L is the linear span of L , L_1 and $[L_1, L_1]$.*

Proof. Since $ad_x : \mathfrak{g} \rightarrow \mathfrak{g}$ is skew symmetric for all $x \in \mathfrak{g}$, it follows that the images of ad_x and $(ad_x)^2$ are equal, hence $[L, L_1] = L_1$.

Let $I := \text{span}(L, L_1, [L_1, L_1]) \subset \mathfrak{g}$. Then $Q([I^\perp, L], \mathfrak{g}) = Q(I^\perp, [L, \mathfrak{g}]) = Q(I^\perp, L_1) = 0$, so that $[I^\perp, L] = 0$. Next, $[I^\perp, L_1] = [I^\perp, [L, L_1]] = [L, [I^\perp, L_1]] \subset [L, \mathfrak{g}] = L_1$. On the other hand, $Q([I^\perp, L_1], L_1) = Q(I^\perp, [L_1, L_1]) = 0$, hence $[I^\perp, L_1] = 0$. Then the Jacobi identity implies that $[I^\perp, [L_1, L_1]] = 0$ as well.

Thus, $[I, I^\perp] = 0$, hence $Q([I, \mathfrak{g}], I^\perp) = Q(\mathfrak{g}, [I, I^\perp]) = 0$, which shows that $I \triangleleft \mathfrak{g}$ is an ideal. On the other hand, any ideal containing L must also contain I which shows the claim. $\blacksquare$

Lemma 4.5 *Let G / K be a compact irreducible symmetric space with Lie algebra $\mathfrak{g}$. Let $\mathfrak{s} \subsetneq \mathfrak{g}$ be a proper subalgebra, and let $\mathfrak{l}_1, \mathfrak{l}_2 \subset \mathfrak{s}$ be subalgebras such that $(ad_{\mathfrak{l}_2} \circ ad_{\mathfrak{l}_1})|_{\mathfrak{s}^\perp}$ vanishes. If there is an $ad_{\mathfrak{s}}$ -invariant and τ -invariant subspace $0 \neq W \subset \mathfrak{s}^\perp$ such that $[\mathfrak{l}_1, W] = W$, then $\mathfrak{l}_2 = 0$.*

Proof. First observe that $[\mathfrak{l}_2, W] = [\mathfrak{l}_2, [\mathfrak{l}_1, W]] \subset (ad_{\mathfrak{l}_2} \circ ad_{\mathfrak{l}_1})(\mathfrak{s}^\perp) = 0$, i.e., $[\mathfrak{l}_2, W] = 0$.

Next, we wish to show that $[\mathfrak{l}_2, [W, \mathfrak{g}]] = 0$. For this, we observe that $[\mathfrak{l}_2, [W, \mathfrak{s}]] = [\mathfrak{l}_2, W] = 0$, and $[\mathfrak{l}_2, [W, W]] = 0$ by the Jacobi identity. Thus, it remains to show that $[\mathfrak{l}_2, [W, W']] = 0$ where $W' := W^\perp \cap \mathfrak{s}^\perp$. Since $Q([W, W'], \mathfrak{s}) = Q(W', [\mathfrak{s}, W]) \subset Q(W', W) = 0$, it follows that $[W, W'] \subset \mathfrak{s}^\perp$. Thus,

$$\begin{aligned} [\mathfrak{l}_2, [W, W']] &= [\mathfrak{l}_2, [[\mathfrak{l}_1, W], W']] \\ &\subset \underbrace{[\mathfrak{l}_2, [\mathfrak{l}_1, [W, W']]]}_{\subset (ad_{\mathfrak{l}_2} \circ ad_{\mathfrak{l}_1})(\mathfrak{s}^\perp) = 0} + [\mathfrak{l}_2, [W, [\mathfrak{l}_1, W']]] \\ &\subset \underbrace{[[\mathfrak{l}_2, W], [\mathfrak{l}_1, W']]}_{=0} + [W, \underbrace{[\mathfrak{l}_2, [\mathfrak{l}_1, W']]}_{\subset (ad_{\mathfrak{l}_2} \circ ad_{\mathfrak{l}_1})(\mathfrak{s}^\perp) = 0}] = 0. \end{aligned}$$

Therefore, $[l_2, W] = [l_2, [W, \mathfrak{g}]] = 0$, and hence, according to Lemma 4.4, $[l_2, I] = 0$ where $I \triangleleft \mathfrak{g}$ is the ideal generated by W . Since $W \neq 0$ is τ -invariant, so is $I \neq 0$, hence the irreducibility of G/K implies that $I = \mathfrak{g}$, i.e., $l_2 \subset \mathfrak{z}(\mathfrak{g}) = 0$. $\blacksquare$

Suppose that $M = H \backslash G/K$ is a minimal biquotient with non-central flats, so that G/K is an irreducible symmetric space by Proposition 4.3, and $\mathcal{F} \neq 0$. Let

$$S := \text{Stab}(\mathcal{F}) = \{g \in G \mid \text{Ad}_g|_{\mathcal{F}} = \text{Id}_{\mathcal{F}}\} \subsetneq G.$$

Evidently, S is compact and τ -invariant, and the Lie algebra $\mathfrak{s}$ of S is the centralizer of $\mathcal{F}$, i.e.,

$$\mathfrak{s} = \{v \in \mathfrak{g} \mid [v, \mathcal{F}] = 0\} \subsetneq \mathfrak{g}.$$

Then $\mathcal{H} \subset \mathfrak{s}$ by (6), and we let $\mathfrak{l} \subset \mathfrak{s}$ be the Lie subalgebra generated by $\mathcal{H}$. Since $\mathcal{H}$ is τ -invariant w.r.t. the involution $\tau : \mathfrak{g} \rightarrow \mathfrak{g}$ from (3), so is $\mathfrak{l}$. Evidently, $\mathcal{F} = \mathfrak{z}(\mathfrak{l})$, hence $\mathfrak{l} = \mathcal{F} \oplus \mathfrak{l}_s$ where $\mathfrak{l}_s$ is semi-simple, and $(\mathfrak{l}_s, \tau)$ is a symmetric pair without Euclidean factor.

If $\mathfrak{l}_s = 0$, i.e., $\mathcal{H} = \mathcal{F}$ is abelian, then $\mathcal{F} \subset \mathfrak{z}(\mathfrak{g})$ by Theorem 3.1. Thus, for biquotients with non-central flats, we may assume that $\mathfrak{l}_s \neq 0$.

Proposition 4.6 *Let $H \backslash G/K$ be a minimal biquotient with non-central flats. Let $\mathfrak{l}_s = \mathfrak{l}_1 \oplus \mathfrak{l}_2$ be a decomposition into τ -invariant ideals such that*

$$\mathcal{H} = \mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_1) \oplus (\mathcal{H} \cap \mathfrak{l}_2). \quad (7)$$

Then either $\mathfrak{l}_k = 0$ or $[\mathfrak{l}_k, \mathfrak{s}^\perp] = \mathfrak{s}^\perp$ for $k = 1, 2$. In particular, $[\mathfrak{l}_s, \mathfrak{s}^\perp] = \mathfrak{s}^\perp$.

Proof. Let us assume that, say, $\mathfrak{l}_1 \neq 0$. Since $\mathfrak{l}_1 \subset \mathfrak{l} \subset \mathfrak{s}$, it is evident that $[\mathfrak{l}_1, \mathfrak{s}^\perp] \subset \mathfrak{s}^\perp$.

Let $G' := \{g \in G \mid \text{Ad}_g(\mathfrak{l}_1 \cap \mathfrak{k}^\perp) = \mathfrak{l}_1 \cap \mathfrak{k}^\perp\}$. Evidently, $G' \subset G$ is a closed subgroup, hence also compact. Since G' and hence its Lie algebra $\mathfrak{g}'$ are τ -invariant, we have the decomposition $\mathfrak{g}' = (\mathfrak{g}' \cap \mathfrak{k}) \oplus (\mathfrak{g}' \cap \mathfrak{k}^\perp) =: \mathfrak{g}'_0 \oplus \mathfrak{g}'_1$, where

$$\mathfrak{g}'_0 = \{k \in \mathfrak{k} \mid [k, \mathfrak{l}_1] \subset \mathfrak{l}_1\}, \quad \text{and} \quad \mathfrak{g}'_1 = \{x \in \mathfrak{k}^\perp \mid [x, \mathfrak{l}_1] = 0\},$$

as $\mathfrak{l}_1$ is generated by $\mathfrak{l}_1 \cap \mathfrak{k}^\perp$. Since $\mathfrak{l}_1$ is semi-simple, we have

$$Q(\mathfrak{g}'_1, \mathfrak{l}_1) = Q(\mathfrak{g}'_1, [\mathfrak{l}_1, \mathfrak{l}_1]) = Q(\underbrace{[\mathfrak{g}'_1, \mathfrak{l}_1]}_{=0}, \mathfrak{l}_1) = 0. \quad (8)$$

Since $\mathcal{F} \oplus \mathfrak{l}_2 \subset \mathfrak{g}'$, we conclude by (7) and (8) that

$$\mathfrak{g}' = \mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_2) \oplus (\mathfrak{g}' \cap \mathcal{H}^\perp) = \mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_2) \oplus (\mathfrak{g}' \cap (\mathfrak{h} \oplus \mathfrak{k})). \quad (9)$$

Let $x \in \text{pr}_{\mathfrak{k}^\perp}(\mathfrak{g}' \cap (\mathfrak{h} \oplus \mathfrak{k}))$. Then $x \in \mathfrak{g}'_1$ and there is a $k \in \mathfrak{k}$ such that $x + k \in \mathfrak{h}$. We have on the one hand $[x + k, \mathcal{H} \cap \mathfrak{l}_1] \subset [\mathfrak{h}, \mathcal{H}] \subset [\mathfrak{h}, \mathfrak{h}^\perp] \subset \mathfrak{h}^\perp$. On the other hand, $[x, \mathcal{H} \cap \mathfrak{l}_1] \subset [\mathfrak{g}'_1, \mathfrak{l}_1] = 0$, hence $[x + k, \mathcal{H} \cap \mathfrak{l}_1] = [k, \mathcal{H} \cap \mathfrak{l}_1] \subset [\mathfrak{k}, \mathfrak{k}^\perp] \subset \mathfrak{k}^\perp$.

Thus, $[x + k, \mathcal{H} \cap \mathfrak{l}_1] = [k, \mathcal{H} \cap \mathfrak{l}_1] \subset \mathfrak{h}^\perp \cap \mathfrak{k}^\perp = \mathcal{H}$, and $Q([k, \mathcal{H} \cap \mathfrak{l}_1], \mathcal{F} \oplus \mathfrak{l}_2) = Q(k, [\mathcal{H} \cap \mathfrak{l}_1, \mathcal{F} \oplus \mathfrak{l}_2]) = 0$, i.e., $[k, \mathcal{H} \cap \mathfrak{l}_1] \subset \mathcal{H} \cap \mathfrak{l}_1$. Since $\mathcal{H} \cap \mathfrak{l}_1$ generates $\mathfrak{l}_1$, it follows that $[k, \mathfrak{l}_1] \subset \mathfrak{l}_1$ and hence, $k \in \mathfrak{g}'_0$, i.e., $x + k \in \mathfrak{h} \cap \mathfrak{g}'$.

Therefore, $pr_{\mathfrak{k}^\perp}(\mathfrak{g}' \cap (\mathfrak{h} \oplus \mathfrak{k})) = pr_{\mathfrak{k}^\perp}(\mathfrak{g}' \cap \mathfrak{h})$, so that we can refine (9) to the (not orthogonal) direct sum decomposition

$$\mathfrak{g}' = \mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_2) \oplus (\mathfrak{g}' \cap \mathfrak{h}) \oplus (\mathfrak{g}' \cap \mathfrak{k}).$$

Thus, for the biquotient $(G' \cap H) \backslash G' / (G' \cap K)$, the horizontal space at $e \in G'$ is $\mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_2) \subsetneq \mathcal{H}$ as $\mathfrak{l}_1 \neq 0$, i.e., $\dim(G' \cap H) \backslash G' / (G' \cap K) < \dim H \backslash G / K$, and $\dim G' \leq \dim G$. Therefore, the minimality of $H \backslash G / K$ implies that $(G' \cap H) \backslash G' / (G' \cap K)$ cannot have non-central flats. Since the centralizer of $\mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_2)$ equals $\mathcal{F}$, this implies that $\mathcal{F} \subset \mathfrak{z}(\mathfrak{g}')$ or, equivalently, $\mathfrak{g}' \subset \mathfrak{s}$.

Let $v \in \mathfrak{s}^\perp \cap [\mathfrak{l}_1, \mathfrak{s}^\perp]^\perp$. Then

$$Q(\underbrace{[v, \mathfrak{l}_1]}_{\subset \mathfrak{s}^\perp}, \mathfrak{s}^\perp) = Q(v, [\mathfrak{l}_1, \mathfrak{s}^\perp]) = 0,$$

hence $[v, \mathfrak{l}_1] = 0$, i.e., $v \in \mathfrak{g}' \subset \mathfrak{s}$. This implies that $v = 0$, i.e. $[\mathfrak{l}_1, \mathfrak{s}^\perp] = \mathfrak{s}^\perp$, which shows our assertion. $\blacksquare$

Let us now investigate the structure of the Lie algebra $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{s}^\perp$ in some more detail. We fix once and for all a *generic* element $v_0 \in \mathcal{F}$, meaning that $[x, \mathcal{F}] = 0$ iff $[x, v_0] = 0$. Thus, the skew-symmetric map $ad_{v_0}|_{\mathfrak{s}^\perp} : \mathfrak{s}^\perp \rightarrow \mathfrak{s}^\perp$ is invertible, hence there is a unique orthogonal complex structure $J : \mathfrak{s}^\perp \rightarrow \mathfrak{s}^\perp$ and a positive definite symmetric map $P_0 : \mathfrak{s}^\perp \rightarrow \mathfrak{s}^\perp$ such that

$$ad_{v_0}|_{\mathfrak{s}^\perp} = JP_0 : \mathfrak{s}^\perp \longrightarrow \mathfrak{s}^\perp. \quad (10)$$

Lemma 4.7 *Let $V_1 := pr_{\mathfrak{s}^\perp}(\mathfrak{h})$ and $V_2 := pr_{\mathfrak{s}^\perp}(\mathfrak{k})$. Then the following hold:*

1. $Q(\mathcal{H}, [V_i, V_i]) = 0$ for $i = 1, 2$, and hence $[\mathcal{H}, V_i] \subset V_i^\perp \cap \mathfrak{s}^\perp$.
2. $\mathfrak{s}^\perp = V_1 \oplus V_2$ and $\dim V_1 = \dim V_2 = \frac{1}{2} \dim \mathfrak{s}^\perp$.
3. As a vector space, $\mathfrak{k} = (\mathfrak{k} \cap \mathfrak{s}) \oplus V_2$.
4. $P_0 V_2 = V_2$ and $V_2 \subset \mathfrak{s}^\perp$ is totally real w.r.t. J , i.e. $JV_2 = V_2^\perp \cap \mathfrak{s}^\perp$.
5. $[\mathfrak{h} \cap \mathfrak{s}, V_1] \subset V_1$.

Proof. Let $x_1, x_2 \in V_1$. Then there exist $k_1, k_2 \in \mathfrak{s} \cap \mathfrak{k}$ such that $x_i + k_i \in \mathfrak{h}$. Thus, for $v \in \mathcal{H}$ it follows

$$\begin{aligned} 0 &= Q(v, \underbrace{[x_1 + k_1, x_2 + k_2]}_{\in \mathfrak{h}}) = Q(v, [x_1, x_2]) + Q(v, \underbrace{[x_1, k_2]}_{\in \mathfrak{s}^\perp} + \underbrace{[k_1, x_2]}_{\in \mathfrak{s}^\perp} + \underbrace{[k_1, k_2]}_{\in \mathfrak{s} \cap \mathfrak{k}}) \\ &= Q(v, [x_1, x_2]), \end{aligned}$$

which shows the first claim for V_1 , and an analogous proof works for V_2 .

If $v_0 \in \mathcal{F}$ is generic, then the form $\omega \in \Lambda^2(\mathfrak{s}^\perp)^*$ defined by $\omega(x, y) := Q(v_0, [x, y]) = Q([v_0, x], y)$ is non-degenerate; indeed, if $0 = \omega(x, \mathfrak{s}^\perp) = Q([v_0, x], \mathfrak{s}^\perp)$ then $[v_0, x] = 0$ which implies that $x = 0$ as v_0 is generic.

Thus, by the first step, $V_i \subset \mathfrak{s}^\perp$ is isotropic w.r.t. ω , hence $\dim V_i \leq \frac{1}{2} \dim \mathfrak{s}^\perp$. Moreover, if $x \in \mathfrak{s}^\perp \cap V_1^\perp \cap V_2^\perp$, then $x \in \mathfrak{h}^\perp \cap \mathfrak{k}^\perp = \mathcal{H} \subset \mathfrak{s}$, so that $x = 0$. This shows that $V_1 + V_2 = \mathfrak{s}^\perp$ and hence the second claim follows.

For the third assertion, suppose there is an $x \in V_2$ and $s_1, s_2 \in \mathfrak{s}$ such that $x + s_1 \in \mathfrak{k}$, $x + s_2 \in \mathfrak{k}^\perp$. Then, since $v_0 \in \mathcal{F} \subset \mathfrak{k}^\perp$ and G/K is symmetric, it follows that $[v_0, x] = [v_0, x + s_i] \in \mathfrak{k}^\perp \cap \mathfrak{k} = 0$, hence $x = 0$. From this, it follows that $\mathfrak{k} = (\mathfrak{k} \cap \mathfrak{s}) \oplus V_2$.

Therefore, $[v_0, [v_0, V_2]] \subset [\mathfrak{k}^\perp, [\mathfrak{k}^\perp, \mathfrak{k}]] \cap \mathfrak{s}^\perp \subset \mathfrak{k} \cap \mathfrak{s}^\perp = V_2$, and by (10) this implies that $P_0^2 V_2 \subset V_2$. But P_0 is symmetric and invertible, hence $P_0 V_2 = V_2$. Thus, $JV_2 = JP_0 V_2 = [v_0, V_2] \subset [\mathfrak{k}^\perp, \mathfrak{k}] \cap \mathfrak{s}^\perp \subset \mathfrak{k}^\perp \cap \mathfrak{s}^\perp = V_2^\perp \cap \mathfrak{s}^\perp$ which shows the fourth part.

For the last assertion, let $h \in \mathfrak{h} \cap \mathfrak{s}$ and $x \in V_1$. Then there is an $s \in \mathfrak{s}$ such that $x + s \in \mathfrak{h}$ and hence $\mathfrak{h} \ni [h, x + s] = [h, x] + [h, s]$. But $[h, s] \subset \mathfrak{s}$, whereas $[h, x] \in \mathfrak{s}^\perp$, i.e., $[h, x] \in pr_{\mathfrak{s}^\perp}(\mathfrak{h}) = V_1$. ■

By virtue of this lemma, we may regard $\mathfrak{s}^\perp \cong V_2 \otimes \mathbb{C}$. We denote the eigenspace decomposition of $\mathfrak{s}^\perp$ w.r.t. P_0 by

$$\mathfrak{s}^\perp = W_1 \oplus \dots \oplus W_n, \quad (11)$$

with eigenvalues $0 < \lambda_1 < \dots < \lambda_n$. Since $v_0 \in \mathfrak{z}(\mathfrak{s}) \cap \mathfrak{k}^\perp$, it follows that W_k is a complex subspace which is invariant both under τ and $ad_{\mathfrak{s}}$. Moreover, since $\mathfrak{s}^\perp = V_1 \oplus V_2$, we have

$$V_1 = \{Jx + P_0^{-1}\phi x \mid x \in V_2\} \quad (12)$$

for some linear map $\phi : V_2 \rightarrow V_2$. By abuse of notation we denote the complex linear extension of ϕ also by $\phi : \mathfrak{s}^\perp \rightarrow \mathfrak{s}^\perp$. Evidently, the latter map is τ -invariant.

Lemma 4.8 *The map $\phi : V_2 \rightarrow V_2$ from (12) is symmetric, and ϕ and P_0 have no common eigenvectors.*

Proof. By Lemma 4.7.1, $Q([v_0, V_1], V_1) = 0$ and thus by (12), we have for $x, y \in V_2$

$$\begin{aligned} 0 &= Q(ad_{v_0}(Jx + P_0^{-1}\phi x), Jy + P_0^{-1}\phi y) \\ &= Q(JP_0(Jx + P_0^{-1}\phi x), Jy + P_0^{-1}\phi y) \\ &= Q(-P_0x + J\phi x, Jy + P_0^{-1}\phi y) \\ &= -Q(P_0x, P_0^{-1}\phi y) + Q(\phi x, y) \\ &= -Q(x, \phi y) + Q(\phi x, y). \end{aligned}$$

For the second part, suppose that $0 \neq x \in V_2$ is such that $P_0x = \lambda x$ and $\phi x = \mu x$. Then $V_1 \ni Jx + P_0^{-1}\phi x = (i + \mu/\lambda)x$, using the identification $\mathfrak{s}^\perp \cong V_2 \otimes \mathbb{C}$, and there is a $k \in \mathfrak{k} \cap \mathfrak{s}$ such that $(i + \mu/\lambda)x + k \in \mathfrak{h}$.

Therefore, $Ad_{\exp t v_0} \mathfrak{h} \ni e^{it\lambda}(i + \mu/\lambda)x + k$. But for a suitable $t_0 \in \mathbb{R}$, we can achieve that $e^{it_0\lambda}(i + \mu/\lambda) \in \mathbb{R}$, i.e., $e^{it_0\lambda}(i + \mu/\lambda)x \in V_2$ and hence $0 \neq e^{it_0\lambda}(i + \mu/\lambda)x + k \in Ad_{\exp t_0 v_0} \mathfrak{h} \cap \mathfrak{k}$ which contradicts the biquotient property. ■

Let $v \in \mathfrak{k}^\perp \cap \mathfrak{s}$. Then $[v, V_2] \subset [\mathfrak{s}, \mathfrak{s}^\perp] \cap [\mathfrak{k}^\perp, \mathfrak{k}] \subset \mathfrak{s}^\perp \cap \mathfrak{k}^\perp = JV_2$, where the last equation follows from Lemma 4.7.4. Therefore, there is a symmetric map $A_v : V_2 \rightarrow V_2$ such that $[A_v, P_0] = 0$ and $ad_v|_{\mathfrak{s}^\perp} = JA_v$.

Lemma 4.9 Let $v \in \mathfrak{k}^\perp \cap \mathfrak{s}$ and let $A_v : V_2 \rightarrow V_2$ be the symmetric linear map such that $ad_v|_{\mathfrak{s}^\perp} = JA_v$.

1. If $v \in \mathcal{H}$ then $[P_0^{-1}A_v, \phi] = 0$.
2. If $v \in \mathcal{V}$ and $x \in \mathfrak{s}^\perp$ is a common eigenvector of $P_0^{-1}A_v$ and ϕ , then $A_v x = 0$.

Proof. Let $v \in \mathcal{H}$ and $x, y \in V_2$, so that $Jx + P_0^{-1}\phi x, Jy + P_0^{-1}\phi y \in V_1$ by (12). By Lemma 4.7.1,

$$\begin{aligned}
0 &= Q([v, Jx + P_0^{-1}\phi x], Jy + P_0^{-1}\phi y) \\
&= Q(JA_v(Jx + P_0^{-1}\phi x), Jy + P_0^{-1}\phi y) \\
&= Q(-A_v x + JA_v P_0^{-1}\phi x, Jy + P_0^{-1}\phi y) \\
&= -Q(A_v x, P_0^{-1}\phi y) + Q(A_v P_0^{-1}\phi x, y) \\
&= -Q(\phi P_0^{-1}A_v x, y) + Q(P_0^{-1}A_v \phi x, y) \quad \text{since } \phi \text{ and } P_0 \text{ are symmetric and } [A_v, P_0] = 0 \\
&= Q([P_0^{-1}A_v, \phi]x, y),
\end{aligned}$$

which shows the first part. For the second, let $v \in \mathcal{V} \cap \mathfrak{k}^\perp \cap \mathfrak{s}$ and $0 \neq x \in V_2$ such that $P_0^{-1}A_v x = \mu x$ and $\phi x = \nu x$. Let $k \in \mathfrak{k} \cap \mathfrak{s}$ be such that $v + k \in \mathfrak{h}$. Thus, $Jx + \nu P_0^{-1}x \in V_1$ by (12), and by Lemma 4.7.5, $(ad_v + ad_k)(V_1) \subset V_1$, so that

$$\begin{aligned}
V_1 &\ni (ad_v + ad_k)(Jx + \nu P_0^{-1}x) \\
&= -A_v x + \nu J P_0^{-1}A_v x + J ad_k x + \nu P_0^{-1}ad_k x \\
&= -\mu P_0 x + \nu \mu Jx + J ad_k x + \nu P_0^{-1}ad_k x \\
&= J(ad_k x + \nu \mu x) + P_0^{-1}(-\mu P_0^2 x + \nu ad_k x).
\end{aligned}$$

Therefore, by (12), $-\mu P_0^2 x + \nu ad_k x = \phi(ad_k x + \nu \mu x)$ or

$$\nu ad_k x - \phi ad_k x = \mu(P_0^2 x + \nu \phi x) = \mu(P_0^2 + \nu^2)x. \quad (13)$$

Since ϕ is symmetric and ad_k is skew-symmetric, we have

$$Q(\nu ad_k x - \phi ad_k x, x) = \nu \underbrace{Q(ad_k x, x)}_{=0} - \underbrace{Q(ad_k x, \overbrace{\phi x}^{\nu x})}_{=0} = 0,$$

and because of (13),

$$\mu Q((P_0^2 + \nu^2)x, x) = 0.$$

Since P_0 and hence $P_0^2 + \nu^2$ is positive definite, this implies that $\mu = 0$, i.e., $A_v x = 0$. ■

We call a subspace $U \subset \mathfrak{s}^\perp$ $(\tau, \phi, \mathfrak{l})$ -invariant if it is invariant under τ , ϕ and $ad_{\mathfrak{l}}$. In particular, U must be ad_{v_0} -invariant, so that each non-zero $(\tau, \phi, \mathfrak{l})$ -invariant subspace is complex and has the form

$$U = U_1 \oplus \dots \oplus U_r, \text{ where } 0 \neq U_k \subset W_{i_k} \text{ for some indices } 1 \leq i_1 < \dots < i_r \leq n$$

with the subspaces W_i from (11). The indices $(i_1, \dots, i_r)$ are called the *weights* of U .

Proposition 4.10 Suppose that $H \backslash G/K$ is a minimal biquotient with non-central flats. Let $0 \neq U \subset \mathfrak{s}^\perp$ be a $(\tau, \phi, \mathfrak{l})$ -invariant subspace. Then $\mathfrak{l}_s$ acts faithfully on U .

Proof. Decompose $\mathfrak{l}_s = \mathfrak{l}_1 \oplus \mathfrak{l}_2$ where $\mathfrak{l}_2 := \text{Ann}(U) \cap \mathfrak{l}_s \triangleleft \mathfrak{l}_s$. We also decompose

$$\mathcal{H} = \mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_1) \oplus (\mathcal{H} \cap \mathfrak{l}_2) \oplus \{v + \alpha(v) \mid v \in \Delta\} \quad (14)$$

for some linear isomorphism $\alpha : \Delta \rightarrow \Delta'$ with $\Delta \subset \mathfrak{l}_1 \cap (\mathcal{H} \cap \mathfrak{l}_1)^\perp$ and $\Delta' \subset \mathfrak{l}_2 \cap (\mathcal{H} \cap \mathfrak{l}_2)^\perp$. Moreover, since U and hence $\mathfrak{l}_i$ are τ -invariant, we have $\Delta, \Delta' \subset \mathfrak{k}^\perp$.

Let $v \in \Delta$. Then $v_1 := v + \alpha(v) \in \mathcal{H}$, whereas $v_2 := v - (\alpha^t)^{-1}v \in \mathcal{V} \cap \mathfrak{k}^\perp$, and $ad_{v_i}|_{\mathfrak{s}^\perp} = JA_i$ for symmetric maps $A_i : V_2 \rightarrow V_2$ for $i = 1, 2$. Since $\alpha(v), (\alpha^t)^{-1}(v) \in \Delta' \subset \text{Ann}(U)$, it follows that $(A_2)|_U = (A_1)|_U$.

But now, Lemma 4.9.1 implies that $[P_0^{-1}A_1, \phi] = 0$, hence $[P_0^{-1}A_2, \phi]|_U = 0$ so that we can find a basis of U consisting of common eigenvectors of ϕ and $P_0^{-1}A_2$. Therefore, Lemma 4.9.2 implies that $A_2(U) = 0$, so that $ad_v(U) = ad_{v_2}(U) = 0$ which implies that $v \in \Delta \cap \mathfrak{l}_2 = 0$. Thus, $\Delta = \Delta' = 0$, i.e. (14) becomes

$$\mathcal{H} = \mathcal{F} \oplus (\mathcal{H} \cap \mathfrak{l}_1) \oplus (\mathcal{H} \cap \mathfrak{l}_2).$$

On the other hand,

$$Q([\mathfrak{l}_2, \mathfrak{s}^\perp], U) = Q(\mathfrak{s}^\perp, \underbrace{[\mathfrak{l}_2, U]}_{=0}) = 0,$$

hence $[\mathfrak{l}_2, \mathfrak{s}^\perp] \subset U^\perp \subsetneq \mathfrak{s}^\perp$, so that Proposition 4.6 implies that $\mathfrak{l}_2 = 0$ which shows the claim. $\blacksquare$

Now define the orthogonal projections $\pi_k : \mathfrak{s}^\perp \rightarrow W_k$ w.r.t. the splitting (11), and the maps

$$\phi_{ij} := \pi_i \circ \phi \circ \pi_j.$$

Since ϕ is symmetric, it follows that $(\phi_{ij})^t = \phi_{ji}$. Also, $\phi = \sum_{i,j} \phi_{ij}$. For a tuple $\sigma = (i_1, \dots, i_n)$ we define

$$\phi_\sigma =: \phi_{i_1 i_2} \circ \phi_{i_2 i_3} \circ \dots \circ \phi_{i_{n-1} i_n},$$

and we say that σ is a tuple from i_1 to i_n . If $i_1 = i_n$ then we call σ an $(i_1\text{-based})$ loop. Note that $(\phi_\sigma)^t = \phi_{\sigma^t}$, where $\sigma^t := (i_n, \dots, i_1)$.

Lemma 4.11 *Let $\sigma = (i_1, \dots, i_n)$ be a tuple from i_1 to i_n and let $v \in \mathcal{H}$. Then*

$$ad_v \circ \phi_\sigma = \frac{\lambda_{i_1}}{\lambda_{i_n}} \phi_\sigma \circ ad_v$$

In particular, $\mathcal{H}$ and hence $\mathfrak{l}$ commutes with ϕ_σ if σ is a loop.

Proof. Evidently, it suffices to show the lemma for tuples $\sigma = (i, j)$. Let $A = A_v : V_2 \rightarrow V_2$ be the symmetric linear map such that $ad_v = JA$. By Lemma 4.9, $[P_0^{-1}A, \phi] = 0$. Now let $A_i := A \circ \pi_i = \pi_i \circ A$. Then $P_0^{-1}A = \sum_i 1/\lambda_i A_i$, hence we have

$$\begin{aligned} 0 &= \pi_i [P_0^{-1}A, \phi] \pi_j \\ &= P_0^{-1}A_i \phi \pi_j - \pi_i \phi P_0^{-1}A_j \\ &= 1/\lambda_i A_i \phi_{ij} - 1/\lambda_j \phi_{ij} A_j \\ &= 1/\lambda_i A \phi_{ij} - 1/\lambda_j \phi_{ij} A, \end{aligned}$$

and from here the claim follows. ■

A $(\tau, \phi, \mathfrak{l})$ -invariant subspace $U \subset V_2$ is called $(\tau, \phi, \mathfrak{l})$ -irreducible if it has no non-trivial $(\tau, \phi, \mathfrak{l})$ -invariant subspace. If $U \subset \mathfrak{s}^\perp$ is $(\tau, \phi, \mathfrak{l})$ -invariant, then so is $\mathfrak{s}^\perp \cap U^\perp$, hence $\mathfrak{s}^\perp$ is the orthogonal sum of $(\tau, \phi, \mathfrak{l})$ -irreducible subspaces.

Definition 4.12 Let $\mathcal{I}$ be the set of all non-zero τ -invariant ideals $\mathfrak{l}' \triangleleft \mathfrak{l}_s$ for which $(\mathfrak{l}', \tau)$ is an irreducible symmetric pair. On $\mathcal{I}$, define the partial ordering $\prec$ by

$$\mathfrak{l}_1 \prec \mathfrak{l}_2 \iff \begin{cases} \text{for all } v \in \mathcal{H}, \text{ we have } pr_{\mathfrak{l}_1}(v) \neq 0 \text{ iff } pr_{\mathfrak{l}_2}(v) \neq 0, \\ \text{and in this case, } \|pr_{\mathfrak{l}_1}(v)\|_{B_{\mathfrak{l}_1}} < \|pr_{\mathfrak{l}_2}(v)\|_{B_{\mathfrak{l}_2}}, \end{cases}$$

where for a compact semi-simple Lie algebra $\mathfrak{l}'$, $\|\cdot\|_{B_{\mathfrak{l}'}}$ denotes the norm w.r.t. the negative of the Killing form.

Note that we have the decomposition $\mathfrak{l}_s := \bigoplus_{\mathfrak{l}' \in \mathcal{I}} \mathfrak{l}'$.

Proposition 4.13 Suppose that $H \backslash G / K$ is a minimal biquotient with non-central flats. Let $0 \neq U = U_1 \oplus \dots \oplus U_r \subset \mathfrak{s}^\perp$ be a $(\tau, \phi, \mathfrak{l})$ -irreducible subspace with weights $i_1 < \dots < i_r$, and let $\mathfrak{l}_{\max}, \mathfrak{l}_{\min} \triangleleft \mathfrak{l}_s$ be maximal and minimal ideals w.r.t. the partial ordering $\prec$. Then

1. $[\mathfrak{l}_{\min}, U_1] = U_1$, and $[\mathfrak{l}_{\min}, U_k] = 0$ for all $k > 1$,
2. $[\mathfrak{l}_{\max}, U_r] = U_r$, and $[\mathfrak{l}_{\max}, U_k] = 0$ for all $k < r$.

Proof. Suppose there is a subspace $0 \neq U'_1 \subset U_1$ which is invariant under τ , $\mathfrak{l}$ and under ϕ_σ for all loops σ . Let $U'_k := \text{span}\{\phi_\sigma(U'_1) \mid \sigma \text{ a tuple from } i_1 \text{ to } i_k\} \subset U_k$. Then U'_k is τ -invariant, and $\phi_{ij}(U'_j) \subset U'_i$ by the definition of ϕ_σ . Also, since for all $v \in \mathcal{H}$, $\phi_\sigma ad_v$ and $ad_v \phi_\sigma$ are multiples of each other by Lemma 4.11, U'_k is also $\mathfrak{l}$ -invariant, so that $\bigoplus_k U'_k \subset U$ is $(\tau, \phi, \mathfrak{l})$ -invariant. But since we assume that U is $(\tau, \phi, \mathfrak{l})$ -irreducible, this means that this direct sum is all of U and hence $U_k = U'_k$ for all k .

Decompose $U_1 := \sum_\alpha U_1^\alpha \otimes \mathbb{R}^{n_\alpha}$ into τ - and $\mathfrak{l}$ -invariant subspaces, such that U_1^α and U_1^β are inequivalent τ -equivariant irreducible representations of $\mathfrak{l}$ for $\alpha \neq \beta$. Since τ and $\mathfrak{l}$ commute with ϕ_σ for all loops σ , it follows that the summands $U_1^\alpha \otimes \mathbb{R}^{n_\alpha}$ are invariant under τ , $\mathfrak{l}$ and these ϕ_σ , so that the preceding paragraph implies that U_1 contains only one such summand. Arguing likewise for all U_k , we conclude $U_k = \tilde{U}_k \otimes \mathbb{R}^{n_k}$ for some irreducible τ -equivariant representation $\tilde{U}_k$ of $\mathfrak{l}$.

For each k , pick a tuple σ_k from i_1 to i_k such that $\phi_{\sigma_k}(U_1) \neq 0$. Then there is an $e_1 \in \mathbb{R}^{n_1}$ such that $(\phi_{\sigma_k})|_{\tilde{U}_1 \otimes e_1} \neq 0$ for all k . Note that $\ker(\phi_{\sigma_k}) = \ker((\phi_{\sigma_k})^t \phi_{\sigma_k}) = \ker \phi_{\sigma_k * \sigma_k^t}$, and since $\sigma_k * \sigma_k^t$ is a loop, it follows that $\ker(\phi_{\sigma_k}) \subsetneq \tilde{U}_1 \otimes e_1$ is a τ - and $\mathfrak{l}$ -invariant proper subspace which must vanish. That is, $\phi_{\sigma_k}|_{\tilde{U}_1 \otimes e_1}$ is injective.

By Lemma 4.11, $ad_v \circ \phi_{\sigma_k}$ and $\phi_{\sigma_k} \circ ad_v$ are multiples of each other for all $v \in \mathcal{H}$, hence $\phi_{\sigma_k}(\tilde{U}_1 \otimes e_1) \subset U_k$ is τ - and $\mathfrak{l}$ -invariant and irreducible, that is, $\phi_{\sigma_k}(\tilde{U}_1 \otimes e_1) = \tilde{U}_k \otimes e_k \subset U_k$ for some $0 \neq e_k \in \mathbb{R}^{n_k}$. Note that e_k is well-defined only up to scalar multiples. However, once we fixed these elements e_k , we may identify $\tilde{U}_k \cong \tilde{U}_1 \otimes e_k$ for $k = 1, \dots, r$ and hence the adjoint representation of $\mathfrak{l}_s$ on U gives rise to a homomorphism

$$ad_{\mathfrak{l}} : \mathfrak{l}_s \longrightarrow \bigoplus_{k=1}^r \text{End}(\tilde{U}_k), \quad (15)$$

and we may regard ϕ_{σ_k} as an isomorphism $\phi_{\sigma_k} : \tilde{U}_1 \rightarrow \tilde{U}_k$.

We decompose the projection of $ad_{\mathfrak{l}}(\mathfrak{l}_s)$ to $End(\tilde{U}_1)$ into irreducible τ -invariant ideals $\mathfrak{u}_1^1 \oplus \dots \oplus \mathfrak{u}_1^m$, and define $\mathfrak{u}_k^\nu := Ad_{\phi_{\sigma_k}}(\mathfrak{u}_1^\nu) \subset End(\tilde{U}_k)$ for $k = 1, \dots, r$ and $\nu = 1, \dots, m$. Once again by Lemma 4.11,

$$Ad_{\phi_{\sigma_k}}^{-1} \left((ad_v)|_{\tilde{U}_k} \right) = (\phi_{\sigma_k}^{-1} \circ ad_v \circ \phi_{\sigma_k})|_{\tilde{U}_1} = \frac{\lambda_{i_1}}{\lambda_{i_k}} (ad_v)|_{\tilde{U}_1}, \quad (16)$$

so that the projection of $ad_{\mathfrak{l}}(\mathfrak{l}_s)$ to $End(\tilde{U}_k)$ equals $\mathfrak{u}_k^1 \oplus \dots \oplus \mathfrak{u}_k^m$. Therefore, we can refine the faithful representation (15) to

$$ad_{\mathfrak{l}} : \mathfrak{l}_s \longrightarrow \bigoplus_{k=1}^r \bigoplus_{\nu=1}^m \mathfrak{u}_k^\nu,$$

where the projection to each summand is surjective. Since $\mathfrak{l}_s$ acts faithfully on U by Proposition 4.10, it follows that $ad_{\mathfrak{l}}^{-1}(\mathfrak{u}_k^\nu) \triangleleft \mathfrak{l}_s$ is an ideal which is τ -equivariantly isomorphic to $\mathfrak{u}_k^\nu$ and hence irreducible, i.e., $ad_{\mathfrak{l}}^{-1}(\mathfrak{u}_k^\nu) \in \mathcal{I}$ with the set $\mathcal{I}$ from Definition 4.12. Thus, we get a map

$$I_0 : \{1, \dots, r\} \times \{1, \dots, m\} \longrightarrow \mathcal{I}, \quad I_0(k, \nu) := ad_{\mathfrak{l}}^{-1}(\mathfrak{u}_k^\nu) \in \mathcal{I}.$$

Let $v \in \mathcal{H}$, and denote by $u_k^\nu \in \mathfrak{u}_k^\nu \subset End(\tilde{U}_k)$ the projection of $(ad_v)|_{\tilde{U}_k}$. Then (16) implies that

$$Ad_{\phi_{\sigma_k}}^{-1}(u_k^\nu) = \frac{\lambda_{i_1}}{\lambda_{i_k}} u_1^\nu. \quad (17)$$

The Killing form is preserved under Lie algebra isomorphisms, and since both $ad_{\mathfrak{l}} : I_0(k, \nu) \rightarrow \mathfrak{u}_k^\nu$ and $Ad_{\phi_{\sigma_k}}^{-1} : \mathfrak{u}_k^\nu \rightarrow \mathfrak{u}_1^\nu$ are isomorphisms, we have

$$\|pr_{I_0(k, \nu)}(v)\|_{B_{I_0(k, \nu)}} = \|u_k^\nu\|_{B_{\mathfrak{u}_k^\nu}} = \|Ad_{\phi_{\sigma_k}}^{-1}(u_k^\nu)\|_{B_{\mathfrak{u}_1^\nu}} = \frac{\lambda_{i_1}}{\lambda_{i_k}} \|u_1^\nu\|_{B_{\mathfrak{u}_1^\nu}},$$

where we used (17) in the last equation. From this, we deduce immediately that

$$\|pr_{I_0(k, \nu)}(v)\|_{B_{I_0(k, \nu)}} = \frac{\lambda_{i_{k'}}}{\lambda_{i_k}} \|pr_{I_0(k', \nu)}(v)\|_{B_{I_0(k', \nu)}},$$

and since for $k < k'$ we have $\lambda_{i_k} < \lambda_{i_{k'}}$, it follows that

$$I_0(k, \nu) \prec I_0(k', \nu) \text{ for all } k < k' \text{ and all } \nu.$$

Thus, if $\mathfrak{l}_{\min} \in \mathcal{I}$ is minimal w.r.t. $\prec$, then $pr_{\mathfrak{u}_k^\nu}(ad_{\mathfrak{l}}(\mathfrak{l}_{\min})) = 0$ for all $k > 1$, that is, $[\mathfrak{l}_{\min}, \tilde{U}_k] = 0$ and therefore $[\mathfrak{l}_{\min}, U_k] = 0$ for all $k > 1$. Since $\mathfrak{l}_{\min} \triangleleft \mathfrak{l}_s$ is a τ -invariant ideal, it follows that $[\mathfrak{l}_{\min}, \tilde{U}_1] \subset \tilde{U}_1$ is τ - and $\mathfrak{l}$ -invariant, hence by irreducibility is either zero or all of $\tilde{U}_1$. In the first case, however, we would have $ad_{\mathfrak{l}_{\min}}|_{s^\perp} = 0$, contradicting the faithfulness of the action of $\mathfrak{l}_s$, hence $[\mathfrak{l}_{\min}, \tilde{U}_1] = \tilde{U}_1$ and therefore $[\mathfrak{l}_{\min}, U_1] = U_1$ as claimed.

The corresponding statements about $\mathfrak{l}_{\max}$ follow analogously. ■

Proposition 4.13 is the last step which is needed to show the

Theorem 4.14 *There are no normal biquotients with non-central flats.*

Proof. Suppose biquotients with non-central flats exist. Then there exists also a *minimal* biquotient with non-central flats $H \backslash G/K$. Let $\mathfrak{l}_{\max}, \mathfrak{l}_{\min} \triangleleft \mathfrak{l}_{\mathfrak{s}} \subset \mathfrak{s}$ and $0 \neq U = U_1 \oplus \dots \oplus U_r \subset \mathfrak{s}^\perp$ be as in Proposition 4.13. If $r = 1$, then any eigenvector of $\phi|_U$ would also be an eigenvector of P_0 as $U \subset W_{i_1}$. But this is impossible by Lemma 4.8.

Thus, $r > 1$ and hence $[\mathfrak{l}_{\max}, [\mathfrak{l}_{\min}, U]] = 0$ by Proposition 4.13. Since $\mathfrak{s}^\perp$ is the direct sum of $(\tau, \phi, \mathfrak{l})$ -irreducible subspaces, it follows that

$$[\mathfrak{l}_{\max}, [\mathfrak{l}_{\min}, \mathfrak{s}^\perp]] = 0.$$

Also, $W_1 = \bigoplus \{U_1 \mid 0 \neq U_1 \oplus \dots \oplus U_r \text{ is } (\tau, \phi, \mathfrak{l})\text{-irreducible with } 1 \text{ as a weight}\}$. Again by Proposition 4.13, this implies that

$$[\mathfrak{l}_{\min}, W_1] = W_1.$$

By Proposition 4.3, G/K is a compact irreducible symmetric space, and if we apply Lemma 4.5 with $\mathfrak{l}_1 := \mathfrak{l}_{\min}$, $\mathfrak{l}_2 := \mathfrak{l}_{\max}$ and $W := W_1$, then we conclude $\mathfrak{l}_{\max} = 0$ which is a contradiction. $\blacksquare$

References

- [E] P.E. Ehrlich, *Metric deformations of curvature, I. Local convex deformations*, Geom.Dedicata **5**, 1–23 (1976)
- [BW] C.Böhm, B.Wilking, *Nonnegatively curved Manifolds with finite fundamental groups admit metrics with positive Ricci curvature*, Geom. Funct. Anal. (to appear)
- [F] W. Fang, *Left invariant metrics on simple compact Lie groups and metrics on spheres*, Ph.D.-thesis, University of Pennsylvania, Philadelphia, USA (2000)
- [N] J.C. Nash, *Positive Ricci curvature on fiber bundles*, J. Differential Geometry **14**, 241–254 (1979)
- [ST] L.J. Schwachhöfer, W. Tuschmann, *Metrics of positive Ricci curvature on quotient spaces*, Math. Ann. **330** (1), 59 – 91 (2004)

FACHBEREICH MATHEMATIK, UNIVERSITÄT DORTMUND, 44221 DORTMUND, GERMANY
 EMAIL: lschwach@math.uni-dortmund.de